arXiv:1601.06310v1 [math.OC] 23 Jan 2016

SOLVING A PROBLEM OF ANGIOGENESIS OF DEGREE THREE

ANASTASIOS N. ZACHOS

Dedicated to Professors Dr. Alexander O. Ivanov and Dr. Alexey A. Tuzhilin for their contributions on minimal trees

Abstract. An absorbing weighted Fermat-Torricelli tree of degree four is a weighted Fermat-Torricelli tree of degree four which is derived as a limiting tree structure from a generalized Gauss tree of degree three (weighted full Steiner tree) of the same boundary convex quadrilateral in $\mathbb{R}^2$. By letting the four variable positive weights which correspond to the fixed vertices of the quadrilateral and satisfy the dynamic plasticity equations of the weighted quadrilateral, we obtain a family of limiting tree structures of generalized Gauss trees which concentrate to the same weighted Fermat-Torricelli tree of degree four (universal absorbing Fermat-Torricelli tree). The values of the residual absorbing rates for each derived weighted Fermat-Torricelli tree of degree four of the universal Fermat-Torricelli tree form a universal absorbing set. The minimum of the universal absorbing Fermat-Torricelli set is responsible for the creation of a generalized Gauss tree of degree three for a boundary convex quadrilateral derived by a weighted Fermat-Torricelli tree of a boundary triangle (Angiogenesis of degree three). Each value from the universal absorbing set contains an evolutionary process of a generalized Gauss tree of degree three.

1. Introduction

We shall describe the structure of a generalized Gauss tree with degree three and a weighted Fermat-Torricelli tree of degree four with respect to a boundary convex quadrilateral $A_1A_2A_3A_4$ in $\mathbb{R}^2$.

Definition 1. [5, Section 2, pp. 2] *A tree topology is a connection matrix specifying which pairs of points from the list $A_1, A_2, A_3, A_4, A_0, A_{0'}$ have a connecting linear segment (edge).*

Definition 2. [2, Subsection 1.2, pp. 8][5],[9] *The degree of a vertex is the number of connections of the vertex with linear segments.*

Let A_1, A_2, A_3, A_4 be four non-collinear points in $\mathbb{R}^2$ and B_i be a positive number (weight) which corresponds to A_i for $i = 1, 2, 3, 4$.

The weighted Fermat-Torricelli problem for four non-collinear points (4wFT problem) in $\mathbb{R}^2$ states that:

1991 *Mathematics Subject Classification.* 51E12, 52A10, 52A55, 51E10.

Key words and phrases. Universal absorbing Fermat-Torricelli set, Universal Fermat-Torricelli minimum value, generalized Gauss problem, weighted Fermat-Torricelli problem, weighted Fermat-Torricelli point, absorbing Fermat-Torricelli tree, absorbing generalized Gauss tree, evolutionary tree.

Problem 1 (4wFT problem)**.** *Find a point (weighted Fermat-Torricelli point)* $A_0 \in \mathbb{R}^2$, *which minimizes*

$$f(A_0) = \sum_{i=1}^{4} B_i \|A_0 - A_i\|, \tag{1.1}$$

where $\|\cdot\|$ *denotes the Euclidean distance*

By letting $B_1 = B_2 = B_3 = B_4$ in the 4wFT problem we obtain the following two cases:

(i) If $A_1A_2A_3A_4$ is a convex quadrilateral, then A_0 is the intersection point of the two diagonals A_1A_3 and A_2A_4,

(ii) If A_i is an interior point of $\triangle A_jA_kA_l$, then $A_0 \equiv A_i$, for $i, j, k, l = 1, 2, 3, 4$ and $i \neq j \neq k \neq l$.

The characterization of the (unique) solution of the 4wFT problem in $\mathbb{R}^2$ is given by the following result which has been proved in [1] and [10]:

Theorem 1. [1, Theorem 18.37, p. 250],[10]

Let A_0 *be a weighted minimum point which minimizes (1.1).*

(a) Then, the 4wFT point A_0 *uniquely exists.*

(b) If for each point $A_i \in \{A_1, A_2, A_3, A_4\}$

$$\| \sum_{j=1, i\neq j}^{4} B_j \vec{u}_{ij} \| > B_i, \tag{1.2}$$

for $i, j = 1, 2, 3, 4$ *holds, then*

(b_1*)* A_0 *does not belong to* $\{A_1, A_2, A_3, A_4\}$ *and*

(b_2*)*

$$\sum_{i=1}^{4} B_i \vec{u}_{0i} = \vec{0}, \tag{1.3}$$

where $\vec{u}_{kl}$ *is the unit vector from* A_k *to* A_l*, for* $k, l \in \{0, 1, 2, 3, 4\}$ *(Weighted Floating Case).*

(c) If there is a point $A_i \in \{A_1, A_2, A_3, A_4\}$ *satisfying*

$$\| \sum_{j=1, i\neq j}^{4} B_j \vec{u}_{ij} \| \leq B_i, \tag{1.4}$$

then $A_0 \equiv A_i$*. (Weighted Absorbed Case).*

The inverse weighted Fermat-Torricelli problem for four non-collinear points (Inverse 4wFT problem) in $\mathbb{R}^2$ states that:

Problem 2. *Inverse 4wFT problem Given a point* A_0 *which belongs to the convex hull of* $A_1A_2A_3A_4$ *in* $\mathbb{R}^2$*, does there exist a unique set of positive weights* B_i*, such that*

$$B_1 + B_2 + B_3 + B_4 = c = const,$$

for which A_0 *minimizes*

$$\sum_{i=1}^{4} B_i \| \|A_0 - A_i\|.$$

By letting $B_4 = 0$ in the inverse 4wFT problem we derive the inverse 3wfT problem which has been introduced and solved by S. Gueron and R. Tessler in [6, Section 4,p. 449].

In 2009, a negative answer with respect to the inverse 4wFT problem is given in [13, Proposition 4.4,p. 417] by deriving a dependence between the four variable weights in $\mathbb{R}^2$. In 2014, we obtain the same dependence of variable weights on some C^2 surfaces in $\mathbb{R}^3$ and we call it the "dynamic plasticity of convex quadrilaterals" ([16, Problem 2, Definition 12, Theorem 1,p.92,p.97-98]).

An important generalization of the Fermat-Torricelli problem is the generalized Gauss problem (or full weighted Steiner tree problem) for convex quadrilaterals in $\mathbb{R}^2$ which has been studied on the K-plane (Sphere, Hyperbolic plane, Euclidean plane) in [15].

We mention the following theorem which provide a characterization for the solutions of the (unweighted)Gauss problem in $\mathbb{R}^2$.

Theorem 2. [1, Theorem(*),pp. 328] *Any solution of the Gauss problem is a Gauss tree (equally weighted full Steiner tree) with at most two (equally weighted) Fermat-Torricelli points (or Steiner points) where each Fermat-Torricelli point has degree three, and the angle between any two edges incident with a Fermat-Torricelli point is of* 120°.

We need to mention all the necessary definitions of the weighted Fermat-Torricelli tree and weighted Gauss tree topologies, in order to derive some important evolutionary structures of the Fermat-Torricelli trees (Absorbing Fermat-Torricelli trees) and Gauss trees (Absorbing Gauss trees) which have been introduced in [17, Definitions 1-7,p. 1070-1071].

Definition 3. *A weighted Fermat-Torricelli tree topology of degree three is a tree topology with all boundary vertices of a triangle having degree one and one interior vertex (weighted Fermat-Torricelli point) having degree three.*

Definition 4. *A weighted Fermat-Torricelli tree topology of degree four is a tree topology with all boundary vertices of a convex quadrilateral having degree one and one interior vertex (4wFT point) having degree four.*

Definition 5. [5, Subsection 3.7,pp. 6] *A weighted Gauss tree topology (or full Steiner tree topology) of degree three is a tree topology with all boundary vertices of a convex quadrilateral having degree one and two interior vertices (weighted Fermat-Torricelli points) having degree three.*

Definition 6. *A weighted Fermat-Torricelli tree of weighted minimum length with a Fermat-Torricelli tree topology of degree four is called a weighted Fermat-Torricelli tree of degree four.*

Definition 7. [5, Subsection 3.7,pp. 6],[17] *A weighted Gauss tree of weighted minimum length with a Gauss tree topology of degree three is called a generalized Gauss tree of degree three or a full weighted minimal Steiner tree.*

In 2014, we study an important generalization of the weighted Gauss (tree) problem that we call a generalized Gauss problem for convex quadrilaterals in $\mathbb{R}^2$ by using a mechanical construction which extends the mechanical construction of Gueron-Tessler in the sense of Pólya and Varigon ([17, Problem 1, Theorem 4, pp.1073-1075]).

We state a generalized Gauss problem for a weighted convex quadrilateral $A_1A_2A_3A_4$. in $\mathbb{R}^2$, such that the weights B_i which correspond to A_i and $B_{00'} \equiv x_G$, satisfy the inequalities

$$|B_i - B_j| < B_k < B_i + B_j,$$

and

$$|B_t - B_m| < B_n < B_t + B_m$$

where x_G is the variable weight which corresponds to the given distance $l \equiv \|A_0 - A_{0'}\|$, for $i, j, k \in \{1, 4, 00'\}$, $t, m, n \in \{2, 3, 00'\}$ and $i \neq j \neq k$, $t \neq m \neq n$.

Problem 3. [17, Problem 1,p. 1073] *Given l, B_1, B_2, B_3, B_4, find a generalized Gauss tree of degree three with respect to $A_1A_2A_3A_4$ which minimizes*

$$B_1\|A_1 - A_0\| + B_2\|A_2 - A_0\| + B_3\|A_3 - A_{0'}\| + B_4\|A_4 - A_{0'}\| + x_G l. \quad (1.5)$$

For $l = 0$, we obtain a weighted Fermat-Torricelli tree of degree four.

Definition 8. [17, Definition 8, p.1076] *We call the variable x_G which depend on l, a generalized Gauss variable.*

Definition 9. [17, Definition 9, p.1080] *The residual absorbing rate of a generalized Gauss tree of degree at most four with respect to a boundary convex quadrilateral is*

$$\sum_{i=1}^{4} B_i - x_G.$$

Definition 10. [17, Definition 10, p.1080] *An absorbing generalized Gauss tree of degree three is a generalized Gauss tree of degree three with residual absorbing rate*

$$\sum_{i=1}^{4} B_i - x_G.$$

In this paper, we prove that the weighted Fermat-Torricelli problem for convex quadrilaterals cannot be solved analytically, by extending the geometrical solution of E. Torricelli and E.Engelbrecht for convex quadrilaterals in $\mathbb{R}^2$ (Section 2, Theorems 3,4)

In section 3, we derive a solution for the generalized Gauss problem in $\mathbb{R}^2$, in the spirit of K. Menger which depends only on five given positive weighta and five Euclidean distances which determine a convex quadrilateral (Section 3, Theorem 5).

In section 4, We give a new approach concerning the dynamic plasticity of quadrilaterals by deriving a new system of equations different from the dynamic plasticity equations which have been deduced in [13, Proposition 4.4,p. 417] and [16, Problem 2, Definition 12, Theorem 1,p.92,p.97-98] (Section 4, Proposition 2). Furthermore, we obtain a surprising connection of the dynamic plasticity of quadrilaterals with a problem of Rene Descarted posed in 1638.

In section 5, we introduce an absorbing weighted Fermat-Torricelli tree of degree four which is derived as a limiting tree structure from a generalized Gauss tree of degree three of the same boundary convex quadrilateral in $\mathbb{R}^2$.

Then, by assuming that the four variable positive weights which correspond to the fixed vertices of the boundary quadrilateral and satisfy the dynamic plasticity

equations, we obtain a family of limiting tree structures of generalized Gauss trees of degree three which concentrate to the same weighted Fermat-Torricelli tree of degree four in a geometric sense (Universal absorbing Fermat-Torricelli tree).

Furthermore, we calculate the values of the universal rates of a Universal absorbing tree regarding a fixed boundary quadrilaterals (Section 5, Examples 2,3).

In section 6, we introduce a class of Euclidean minimal tree structures that we call steady trees and evolutionary trees (Section 6, Definitions 15,16). Thus, the minimum of the universal absorbing Fermat-Torricelli set (Universal Fermat-Torricelli minimum value) leads to the creation of a generalized Gauss tree of degree three for the same boundary convex quadrilateral which is derived by a weighted Fermat-Torricelli tree of degree four. A universal absorbing Fermat-Torricelli minimum value corresponds to the intersection point (4wFT point). This quantity is of fundamental importance, because by attaining this value the absorbing Fermat-Torricelli tree start to grow and will be able to produce a generalized Gauss tree of degree three (Evolutionary tree). Each specific value from the universal absorbing set gives an evolutionary process of a generalized Gauss tree of degree three regarding a fixed boundary quadrilateral by spending a positive quantity from the storage of the universal Fermat-Torricelli quantity which stimulates the evolution at the 4wFT point (Section 6, Example 4, Angiogenesis of degree three).

2. Extending Torricelli-Engelbrecht's solution for convex quadrilaterals

The weighted Torricelli-Engelbrecht solution for a triangle $\triangle A_1A_2A_3$ in the weighted floating case is given by the following proposition:

Lemma 1. [6],[4] *If A_0 is an interior weighted Fermat-Torricelli point of $\triangle A_1A_2A_3$, then*

$$\angle A_iA_0A_j \equiv \alpha_{i0j} = \arccos\left(\frac{B_k^2 - B_i^2 - B_j^2}{2B_iB_j}\right). \tag{2.1}$$

Let $A_1A_2A_3A_4$ be a convex quadrilateral in $\mathbb{R}^2$, O be the intersection point of the two diagonals and B_i be a given weight which corresponds to the vertex A_i, A_0 be the weighted Fermat-Torricelli point in the weighted floating case (Theorem) and $\vec{U}_{ij}$ be the unit vector from A_i to A_j, for $i, j = 1, 2, 3, 4$.

We mention the geometric plasticity principle of quadrilaterals in $\mathbb{R}^2$,

Lemma 2. [13],[16, Definition 13,Theorem 3, Proposition 8, Corollary 4 p. 103-108] *Suppose that the weighted floating case of the weighted Fermat-Torricelli point A_0 point with respect to $A_1A_2A_3A_4$ is satisfied:*

$$\left\|B_i\vec{U}_{ki} + B_j\vec{U}_{kj} + B_m\vec{U}_{km}\right\| > B_k,$$

for each $i, j, k, m = 1, 2, 3, 4$ and $i \neq j \neq k \neq m$. If A_0 is connected with every vertex A_k for $k = 1, 2, 3, 4$ and we select a point A'_k with non-negative weight B_k which lies on the ray A_kA_0 and the quadrilateral $A'_1A'_2A'_3A'_4$ is constructed such that:

$$\left\|B_i\vec{U}_{k'i'} + B_j\vec{U}_{k'j'} + B_m\vec{U}_{k'm'}\right\| > B_k,$$

for each $i', j', k', m' = 1, 2, 3, 4$ and $i' \neq j' \neq k' \neq m'$. Then the weighted Fermat-Torricelli point A'_0 is identical with A_0.

Theorem 3. *The weighted Torricelli-Engelbrecht solution for $A_1A_2A_3A_4$ is given by the following system of four equations w.r. to the variables α_{102}, α_{203}, α_{304} and α_{401} :*

$$\csc^2\alpha_{102}\csc^2\alpha_{304}\csc^2\alpha_{401}\left(\cos\alpha_{102}-\sin\alpha_{102}\right)\left(\cos\alpha_{304}-\sin\alpha_{304}\right)$$
$$\left(\cos\left(\alpha_{102}-\alpha_{304}\right)-\cos\left(\alpha_{102}+\alpha_{304}+2\alpha_{401}\right)-2\sin\left(\alpha_{102}+\alpha_{304}\right)\right)=0, \quad (2.2)$$

$$-B_1^2-2B_1B_2\cos\alpha_{102}-B_2^2+B_3^2-2B_1B_4\cos\alpha_{401}-2B_2B_4\cos\left(\alpha_{102}+\alpha_{401}\right)-B_4^2=0, \quad (2.3)$$

$$\alpha_{304}=\arccos\left(\frac{B_1^2+2B_1B_2\cos\alpha_{102}+B_2^2-B_3^2-B_4^2}{2B_3B_4}\right) \quad (2.4)$$

and

$$\alpha_{203}=2\pi-\alpha_{102}-\alpha_{304}-\alpha_{401}. \quad (2.5)$$

Proof. Assume that we select B_1, B_2, B_3, B_4, such that A_0 is an interior point of $\triangle A_1OA_2$. By applying the geometric plasticity principle of Lemma 2 we could choose a transformation of $A_1A_2A_3A_4$ to the square $A_1'A_2'A_3'A_4'$, where $A_0'=A_0$ and A_O is an interior point of $A_1'O'A_2'$ where O' is the intersection of $A_1'A_3'$ and $A_2'A_4'$.

We consider the equations of the three circles which pass through A_1, A_2, A_0, A_1, A_4, A_0 and A_3, A_4, A_0, respectively, which meet at A_0 :

$$\left(x-\frac{a}{2}\right)^2+\left(y-\frac{1}{2}a\cot\alpha_{102}\right)^2=\frac{1}{4}a^2\csc^2\alpha_{102} \quad (2.6)$$

$$\left(y-\frac{a}{2}\right)^2+\left(x-\frac{1}{2}a\cot\alpha_{401}\right)^2=\frac{1}{4}a^2\csc^2\alpha_{401} \quad (2.7)$$

$$\left(x-\frac{a}{2}\right)^2+\left(y-(a-\frac{1}{2}a\cot\alpha_{304})\right)^2=\frac{1}{4}a^2\csc^2\alpha_{304} \quad (2.8)$$

By subtracting (2.7) from (2.6), (2.8) from (2.6) and solving w.r. to x, y we get:

$$x=\frac{a(-1+\cot\alpha_{102})(-1+\cot\alpha_{304})}{(-2+\cot\alpha_{102}+\cot\alpha_{304})(-1+\cot\alpha_{401})} \quad (2.9)$$

and

$$y=-\frac{-a\cot\alpha_{304}+a}{\cot\alpha_{102}+\cot\alpha_{304}-2} \quad (2.10)$$

By substituting (2.9) and (2.10) in (2.6), we obtain (2.2).

Taking into account the weighted floating equilibrium condition, we get:

$$-B_3\vec{u}_{30}=B_1\vec{u}_{10}+B_2\vec{u}_{20}+B_4\vec{u}_{40} \quad (2.11)$$

or

$$B_1\vec{u}_{10}+B_2\vec{u}_{20}=-B_3\vec{u}_{30}-B_4\vec{u}_{40}. \quad (2.12)$$

By squaring both parts of (2.11) we derive (2.3) and by squaring both parts of (2.12) we derive (2.4).



FIGURE 1. A generalized Gauss Menger tree of degree three regarding a boundary convex quadrilateral

□

Remark 1. *A different approach was used in* [13, Solution 2.2,Example 2.4, p. 413-414]*, in order to derive a similar system of equations w.r. to* α_{401} *and* α_{102}.

Example 1. *By substituting* $B_1 = 3.5$, $B_2 = 2.5$, $B_3 = 2$, $B_4 = 1$, $a = 10$ *in (2.2) and (2.3) and solving this system of equations numerically by using for instance Newton method and choosing as initial values* $\alpha^o_{102} = 2.7$ *rad,* $\alpha^o_{401} = 1.2$ *rad, we obtain* $\alpha_{102} = 2.30886$ *and* $\alpha_{401} = 1.57801$ *rad. By substituting* $\alpha_{102} = 2.30886$ *and* $\alpha_{401} = 1.57801$ *rad in (2.4) we get* $\alpha_{304} = 1.12492$ *rad. From (2.5), we get* $\alpha_{203} = 1.2714$ *rad. By substituting the angles* α_{i0j} *in (2.9) and (2.10), we derive* $x = 4.0700893$ *and* $y = 2.146831$.

Theorem 4. *There does not exist an analytical solution for the 4wFT problem in* $\mathbb{R}^2$.

Proof. The system of the two equations (2.2) and (2.3) taking into account (2.4) cannot be solved explicitly w.r to α_{102} and α_{401}. Therefore, by considering (2.9) and (2.10) we deduce that the location of the 4wFT point A_0 cannot also be expressed explicitly via the angles α_{i0j}, for $i, j = 1, 2, 3, 4$, for $i \neq j$. □

Thus, from Theorem 4 the position of a weighted Fermat-Torricelli tree of degree four cannot be expressed analytically and may be found by using numerical methods (see also in [13]).

3. An absorbing generalized Gauss-Menger tree in $\mathbb{R}^2$

Let $A_1A_2A_3A_4$ be a boundary weighted convex quadrilateral of a generalized Gauss tree of degree three in $\mathbb{R}^2$ and A_0, $A_{0'}$ are the two weighted Fermat-Torricelli (3wFT) points of degree three which are located at the convex hull of the boundary quadrilateral.

We denote by $l \equiv \|A_0 - A_{0'}\|$, $a_{ij} \equiv \|A_i - A_j\|$, $\alpha_{ijk} \equiv \angle A_iA_jA_k$, $a_{10} \equiv a_1$, $a_{40} \equiv a_4$, $a_{20'} \equiv a_2$, $a_{30'} \equiv a_3$ (Fig. 1) and by $B_i \equiv \frac{B'_i}{\sum_{i=1}^4 B'_i}$, for $i, j, k \in \{0, 0', 1, 2, 3, 4\}$ and $i \neq j \neq k$.

We proceed by giving the following two lemmas which have been proved recently in [15, Theorem 1] and [14].

Lemma 3. [15, Theorem 1],[14, Theorem 2.1,p. 485]

A generalized Gauss tree of degree three (full weighted Steiner minimal tree) of $A_1A_2A_3A_4$ consists of two weighted Fermat-Torricelli points A_0, A_0' which are located at the interior convex domain with corresponding given weights B_0, $B_{0'}$ and minimizes the objective function:

$$B_1a_1 + B_2a_2 + B_3a_3 + B_4a_4 + \frac{B_0 + B_{0'}}{2}l \to min, \tag{3.1}$$

such that:

$$|B_i - B_j| < B_k < B_i + B_j, \tag{3.2}$$

$$|B_t - B_m| < B_n < B_t + B_m \tag{3.3}$$

where

$$B_{00'} \equiv \frac{B_0 + B_{0'}}{2},$$

for $i, j, k \in \{1, 4, 00'\}$, $t, m, n \in \{2, 3, 00'\}$ and $i \neq j \neq k$, $t \neq m \neq n$,

Suppose that B_1, B_2, B_3, B_4, $B_{00'}$ satisfy the inequalities (3.2) and (3.3).

Lemma 4. [14, Theorem 2.2, p. 486] *The location of A_0 and $A_{0'}$ is given by the relations:*

$$\cot\varphi = \frac{B_0a_{12} + B_4a_{14}\cos(\alpha_{214} - \alpha_{400'}) + B_3a_{23}\cos(\alpha_{123} - \alpha_{30'0})}{B_4a_{14}\sin(\alpha_{214} - \alpha_{400'}) - B_3a_{23}\sin(\alpha_{123} - \alpha_{30'0})}, \tag{3.4}$$

$$a_1 = \frac{a_{14}\sin(\alpha_{214} - \varphi - \alpha_{400'})}{\sin(\alpha_{100'} + \alpha_{400'})} \tag{3.5}$$

and

$$a_2 = \frac{a_{23}\sin(\alpha_{123} + \varphi - \alpha_{300'})}{\sin(\alpha_{20'0} + \alpha_{30'0})} \tag{3.6}$$

where φ is the angle which is formed between the line defined by A_1 and A_2 and the line which passes from A_1 and it is parallel to the line defined by A_0 and A_0'.

Definition 11. *A generalized Gauss-Menger tree is a solution of a generalized Gauss problem in $\mathbb{R}^2$ for a boundary quadrilateral $A_1A_2A_3A_4$ which depend on the Euclidean distances a_{ij} and the five given weights B_1, B_2, B_3, B_4 and $B_{00'}$.*

By letting $B_{00'} \equiv x_G$, we obtain an absorbing generalized Gauss-Menger tree for a boundary quadrilateral.

Theorem 5. *An absorbing generalized Gauss-Menger tree w.r. to a fixed convex quadrilateral $A_1A_2A_3A_4$ depends only on the five given weights B_i, B_2, B_3, B_4, $B_{00'} \equiv x_G$ and the five given lengths a_{12}, a_{23}, a_{34}, a_{41} and a_{13}.*

Proof. Consider the Caley-Menger determinant which gives the volume of a tetrahedron $A_1A_2A_3A_4$ in $\mathbb{R}^3$.

$$288V^2 = \det \begin{pmatrix} 0 & a_{12}^2 & a_{13}^2 & a_{14}^2 & 1 \\ a_{12}^2 & 0 & a_{23}^2 & a_{24}^2 & 1 \\ a_{13}^2 & a_{23}^2 & 0 & a_{34}^2 & 1 \\ a_{14}^2 & a_{24}^2 & a_{34}^2 & 0 & 1 \\ 1 & 1 & 1 & 1 & 0 \end{pmatrix}. \tag{3.7}$$

By letting $V = 0$ in (3.7), we obtain a dependence of the six distances a_{12}, a_{13}, a_{14}, a_{23}, a_{34} and a_{24}. For instance, by solving a fourth order degree equation w.r. to a_{13} we derive that $a_{13} = (a_{12}, a_{14}, a_{23}, a_{34}, a_{24})$.

By applying the cosine law in $\triangle A_1A_2A_4$ and $\triangle A_1A_2A_3$ we get:

$$\alpha_{214} = \arccos\left(\frac{a_{12}^2 + a_{14}^2 - a_{24}^2}{2a_{12}a_{14}}\right), \tag{3.8}$$

and

$$\alpha_{123} = \arccos\left(\frac{a_{12}^2 + a_{23}^2 - a_{13}^2}{2a_{12}a_{23}}\right). \tag{3.9}$$

By Lemma 1 and Lemma 3 and considering that A_0 is the 3wFT point of $\triangle A_1A_4A_{0'}$ and $A_{0'}$ is the 3wFT point of $\triangle A_2A_3A_0$ we get:

$$\alpha_{100'} = \arccos\left(\frac{B_4^2 - B_1^2 - x_G^2}{2B_1x_G}\right), \tag{3.10}$$

$$\alpha_{0'04} = \arccos\left(\frac{B_1^2 - B_4^2 - x_G^2}{2B_4x_G}\right), \tag{3.11}$$

$$\alpha_{104} = \arccos\left(\frac{x_G^2 - B_1^2 - B_4^2}{2B_1B_4}\right), \tag{3.12}$$

$$\alpha_{00'3} = \arccos\left(\frac{B_2^2 - B_3^2 - x_G^2}{2B_3x_G}\right), \tag{3.13}$$

$$\alpha_{00'2} = \arccos\left(\frac{B_3^2 - x_G^2 - B_2^2}{2x_GB_2}\right), \tag{3.14}$$

and

$$\alpha_{20'3} = \arccos\left(\frac{x_G^2 - B_2^2 - B_3^2}{2B_2B_3}\right). \tag{3.15}$$

Therefore, by replacing (3.10), (3.11), (3.14), (3.13), (3.8), (3.9) in (3.4), (3.5), (3.6) and taking into account the dependence of the six distances a_{ij}, for $i, j = 1, 2, 3, 4$, we derive that φ, a_1 and a_2 depend only on B_1, B_2, B_3, B_4, x_G and a_{12}, a_{13}, a_{23}, a_{34}, a_{24}.

□

4. The dynamic plasticity of convex quadrilaterals

In this section, we deal with the solution of the inverse 4wFT problem in $\mathbb{R}^2$ which has been introduced in [13] and developed in [16], in order to obtain a new system of equations of the dynamic plasticity of weighted quadrilaterals w.r. to the four variable weights $(B_i)_{1234}$ for $i = 1, 2, 3, 4$, which cover also the case $(B_1)_{1234} = (B_3)_{1234}$ and $(B_2)_{1234} = (B_4)_{1234}$.

First, we start by mentioning the solution of S. Gueron and R. Tessler ([6, Section 4,p. 449]) of the inverse 3wFT problem for three non-collinear points in $\mathbb{R}^2$. By

letting $B_i = 0$ in the inverse 4wFT problem for convex quadrilaterals (Problem 2) we derive the inverse 3wFT problem for s triangle.

Consider the inverse 3wFT problem for $\triangle A_iA_jA_k$ in $\mathbb{R}^2$.

Lemma 5. [6, Section 4,p. 449] *The unique solution of the inverse 3wFT problem for $\triangle A_iA_jA_k$ is given by*

$$(\frac{B_i}{B_j})_{ijk} = \frac{\sin\alpha_{jik}}{\sin\alpha_{ijk}}. \tag{4.1}$$

Definition 12. [16] *We call dynamic plasticity of a weighted Fermat-Torricelli tree of degree four the set of solutions of the four variable weights with respect to the inverse 4wFT problem in $\mathbb{R}^2$ for a given constant value c which correspond to a family of weighted Fermat-Torricelli tree of degree four that preserve the same Euclidean tree structure (the corresponding 4wFT point remains the same for a fixed boundary convex quadrilateral), such that the three variable weights depend on a fourth variable weight and the value of c.*

By taking into account Lemma 5 for the triangles $\triangle A_1A_2A_3$, $\triangle A_1A_3A_4$, $\triangle A_1A_2A_4$ and the weighted floating equilibrium condition (1.3) taken from Theorem 1 ([13], [16])

Proposition 1. [13, Proposition 4.4,p. 417],[16, Problem 2, Definition 12, Theorem 1,p.92,p.97-98] *Suppose that A_0 does not belong to the intersection of the linear segments A_1A_3 and A_2A_4. The dynamic plasticity of the variable weighted Fermat-Torricelli tree of degree four in $\mathbb{R}^2$ is given by the following three equations:*

$$(\frac{B_2}{B_1})_{1234} = (\frac{B_2}{B_1})_{123}[1 - (\frac{B_4}{B_1})_{1234}(\frac{B_1}{B_4})_{134}], \tag{4.2}$$

$$(\frac{B_3}{B_1})_{1234} = (\frac{B_3}{B_1})_{123}[1 - (\frac{B_4}{B_1})_{1234}(\frac{B_1}{B_4})_{124}], \tag{4.3}$$

and

$$(B_1)_{1234} + (B_2)_{1234} + (B_3)_{1234} + (B_4)_{1234} = c = const. \tag{4.4}$$

It is worth mentioning that each quad of values $\{(B_1)_{1234}, (B_2)_{1234}, (B_3)_{1234}, (B_4)_{1234}\}$, which satisfy simultaneously (4.2), (4.3), (4.4) create a unique concentration of different families of tetrafocal ellipses (Polyellipse or Egglipse) to the same 4wFT point A_0 of $A_1A_2A_3A_4$, A family of tetrafocal ellipses may be constructed by selecting a decreasing sequence of real numbers $c_n((B_1)_{1234}, (B_2)_{1234}, (B_3)_{1234}, (B_4)_{1234}))$

$$c_n((B_1)_{1234}, (B_2)_{1234}, (B_3)_{1234}, (B_4)_{1234}); X) = \sum_{i=1}^{4}(B_i)_{1234}\|A_i - X\|$$

which converge to the constant number

$$c((B_1)_{1234}, (B_2)_{1234}, (B_3)_{1234}, (B_4)_{1234}) \equiv f(A_0, (B_1)_{1234}, (B_2)_{1234}, (B_3)_{1234}, (B_4)_{1234}).$$

or

$$c((B_1)_{1234}, (B_2)_{1234}, (B_3)_{1234}, (B_4)_{1234}) = \sum_{i=1}^{4}(B_i)_{1234}\|A_i - A_0\|$$

or

$$c((B_1)_{1234},(B_2)_{1234},(B_3)_{1234},(B_4)_{1234})=\sum_{i=1}^{4}(B_i)_{1234}a_i.$$

The concentration of different families of tetrafocal ellipses to the same point provide a surprising connection with a problem posed by R. Descartes in 1638. According to [1, Chapter II,p. 235], in a letter from August 23, 1638, R. Descartes invited P. de Fermat to investigate the properties of tetrafocal ellipses in $\mathbb{R}^2$. The dynamic plasticity of weighted quadrilaterals solves the problem of concentration of tetrafocal ellipse and offers a new property to R.Descartes' problem.

We proceed by deriving a new system of dynamic plasticity equations for a weighted Fermat-Torricelli tree of degree four which also includes a class of weighted Fermat-Torricelli trees of degree four which coincides with the two diagonals of the boundary convex quadrilateral for $(B_1)_{1234}=(B_3)_{1234}$ and $(B_2)_{1234}=(B_4)_{1234}$.

Proposition 2. *The dynamic plasticity of the variable weighted Fermat-Torricelli tree of degree four in $\mathbb{R}^2$ is given by the following three equations:*

$$\begin{aligned}(B_1)_{1234}^2+(B_2)_{1234}^2+2(B_1)_{1234}(B_2)_{1234}\cos\alpha_{102}=\\ (B_3)_{1234}^2+(B_4)_{1234}^2+2(B_3)_{1234}(B_4)_{1234}\cos\alpha_{304},\end{aligned}\tag{4.5}$$

$$\begin{aligned}(B_1)_{1234}^2+(B_4)_{1234}^2+2(B_1)_{1234}(B_4)_{1234}\cos\alpha_{104}=\\ (B_2)_{1234}^2+(B_3)_{1234}^2+2(B_2)_{1234}(B_3)_{1234}\cos\alpha_{203},\end{aligned}\tag{4.6}$$

and

$$(B_1)_{1234}+(B_2)_{1234}+(B_3)_{1234}+(B_4)_{1234}=cconst.\tag{4.7}$$

Proof. Suppose that we select four positive weights $(B_i)_{1234}(0)$ which correspond the vertex A_i of the boundary convex quadrilateral $A_1A_2A_3A_4$, such that the weighted floating inequalities (1.2) of Theorem 1 hold, in order to locate the 4wFT point A_0 at the interior of $A_1A_2A_3A_4$.

From the weighted floating (variable weighted) equilibrium condition of the 4wFT point (1.3)we get

$$(B_1)_{1234}\vec{u}_{10}+(B_2)_{1234}\vec{u}_{20}=-(B_3)_{1234}\vec{u}_{30}-(B_4)_{1234}\vec{u}_{40}\tag{4.8}$$

and

$$(B_1)_{1234}\vec{u}_{10}+(B_3)_{1234}\vec{u}_{30}=-(B_2)_{1234}\vec{u}_{20}-(B_4)_{1234}\vec{u}_{40},\tag{4.9}$$

which yield (4.5) and (4.6), respectively.

□

Corollary 1. *If the variable weighted 4wFT point is the intersection point of A_1A_3 and A_2A_4, then*

$$(B_1)_{1234}=(B_3)_{1234}\tag{4.10}$$

and

$$(B_2)_{1234}=(B_4)_{1234}.\tag{4.11}$$

$(B_1)_{1234}=(B_3)_{1234}$ *and* $(B_2)_{1234}=(B_4)_{1234}$.

Proof. By letting $\alpha_{102}=\alpha_{304}$ and $\alpha_{104}=\alpha_{203}$, in (4.5) and (4.6) we obtain (4.10) and (4.11), respectively. □

Remark 2. *The dynamic plasticity equations of Theorem 1 depend on the solutions of the inverse 3FT problem for $\triangle A_1A_2A_3$, $\triangle A_1A_3A_4$ and $\triangle A_1A_2A_4$. Thus, the corresponding 4wFT point A_0 remains at the interior of $\triangle A_1A_2A_3$, in order to derive from the inverse 3wFT problem for $\triangle A_1A_2A_3$ the inverse wFT problem for $A_1A_2A_3A_4$. The dynamic plasticity equations of Theorem 2 generalizes the dynamic plasticity equations of Theorem 1 because A_0 could also lie on the side A_1A_3 of $\triangle A_1A_2A_3$ and the side A_2A_4 of $\triangle A_1A_2A_4$. These are the cases where the inverse 3wFT problem for $\triangle A_1A_2A_3$ and $\triangle A_1A_2A_4$ do not hold.*

5. A universal Fermat-Torricelli minimal value of a family of absorbing generalized Gauss trees of degree three

Suppose that an absorbing weighted Fermat-Torricelli tree of degree four is derived as a limiting tree structure from an absorbing generalized Gauss-Menger tree of degree three regarding a fixed boundary quadrilateral $A_1A_2A_3A_4$ for a specific value c_G of the generalized Gauss variable.

We need to consider the following lemma which gives $\bar{B}_i$ as a linear function of $\bar{B}_4$ regarding a fixed variable weighted Fermat-Torricelli tree of degree four in $\mathbb{R}^2$.

Lemma 6. [13, Corollary 4.5,p. 418]*Let* $\sum_{1234}\bar{B}:=(\bar{B}_1)_{1234}(1+\frac{\bar{B}_2}{\bar{B}_1}+\frac{\bar{B}_3}{\bar{B}_1}+\frac{\bar{B}_4}{\bar{B}_1})_{1234}$. *If* $\sum_{1234}\bar{B}=\sum_{123}\bar{B}=\sum_{124}\bar{B}=\sum_{134}\bar{B}$, *then*

$$(\bar{B}_i)_{1234}=x_i(\bar{B}_4)_{1234}+y_i, i=1,2,3:$$

$$(x_1,y_1)=(\frac{(\frac{\bar{B}_1}{\bar{B}_4})_{134}(\frac{\bar{B}_2}{\bar{B}_1})_{123}+(\frac{\bar{B}_1}{\bar{B}_4})_{124}(\frac{\bar{B}_3}{\bar{B}_1})_{123}-1}{1+(\frac{\bar{B}_2}{\bar{B}_1})_{123}+(\frac{\bar{B}_3}{\bar{B}_1})_{123}},(\bar{B}_1)_{123})$$

$$(x_2,y_2)=(x_1(\frac{\bar{B}_2}{\bar{B}_1})_{123}-(\frac{\bar{B}_1}{\bar{B}_4})_{134}(\frac{\bar{B}_2}{\bar{B}_1})_{123},(\bar{B}_2)_{123})$$

$$(x_3,y_3)=(x_1(\frac{\bar{B}_3}{\bar{B}_1})_{123}-(\frac{\bar{B}_1}{\bar{B}_4})_{124}(\frac{\bar{B}_3}{\bar{B}_1})_{123},(\bar{B}_3)_{123}).$$

Theorem 6. *A universal constant does not exist for a unique fixed (variable) weighted Fermat-Torricelli tree of degree four which is obtained as a limiting tree structure from a family of variable weighted Gauss-Menger trees (or full variable weighted Steiner trees) w.r. to a fixed boundary convex quadrilateral $A_1A_2A_3A_4$, which depend on the variable weights $(\bar{B}_i)_{1234}$ which satisfy the dynamic plasticity equations*

$$(\frac{\bar{B}_2}{\bar{B}_1})_{1234}=(\frac{\bar{B}_2}{\bar{B}_1})_{123}[1-(\frac{\bar{B}_4}{\bar{B}_1})_{1234}(\frac{\bar{B}_1}{\bar{B}_4})_{134}],\tag{5.1}$$

$$(\frac{\bar{B}_3}{\bar{B}_1})_{1234}=(\frac{\bar{B}_3}{\bar{B}_1})_{123}[1-(\frac{\bar{B}_4}{\bar{B}_1})_{1234}(\frac{\bar{B}_1}{\bar{B}_4})_{124}],\tag{5.2}$$

and

$$(\bar{B}_1)_{1234}+(\bar{B}_2)_{1234}+(\bar{B}_3)_{1234}+(\bar{B}_4)_{1234}=cconstant.\tag{5.3}$$

or

$$(\bar{B}_1)^2_{1234} + (\bar{B}_2)^2_{1234} + 2(\bar{B}_1)_{1234}(\bar{B}_2)_{1234}\cos\alpha_{102} = (\bar{B}_3)^2_{1234} + (\bar{B}_4)^2_{1234} + 2(\bar{B}_3)_{1234}(\bar{B}_4)_{1234}\cos\alpha_{304}, \tag{5.4}$$

$$(\bar{B}_1)^2_{1234} + (\bar{B}_4)^2_{1234} + 2(\bar{B}_1)_{1234}(\bar{B}_4)_{1234}\cos\alpha_{104} = (\bar{B}_2)^2_{1234} + (\bar{B}_3)^2_{1234} + 2(\bar{B}_2)_{1234}(\bar{B}_3)_{1234}\cos\alpha_{203}, \tag{5.5}$$

and

$$(\bar{B}_1)_{1234} + (\bar{B}_2)_{1234} + (\bar{B}_3)_{1234} + (\bar{B}_4)_{1234} = cconst. \tag{5.6}$$

Proof. Suppose that we select four positive weights $(\bar{B}_i)_{1234}(0)$ which correspond the vertex A_i of the boundary convex quadrilateral $A_1A_2A_3A_4$, such that the weighted floating inequalities (1.2) of Theorem 1 hold, in order to locate the 4wFT point A_0 at the interior of $A_1A_2A_3A_4$ and particularly at the interior of $\triangle A_1OA_2$, where O is the intersection of the diagonals A_1A_3 and A_2A_4, closer to the vertex A_1.

The location of the 3wFT points A_0 and $A_{0'}$, respectively, is given by the relations (Lemma 4):

$$\cot\varphi = \frac{x_G a_{12} + \bar{B}_4 a_{14}\cos(\alpha_{214} - \alpha_{400'}) + \bar{B}_3 a_{23}\cos(\alpha_{123} - \alpha_{30'0})}{\bar{B}_4 a_{14}\sin(\alpha_{214} - \alpha_{400'}) - \bar{B}_3 a_{23}\sin(\alpha_{123} - \alpha_{30'0})}, \tag{5.7}$$

$$a_1 = \frac{a_{14}\sin(\alpha_{214} - \varphi - \alpha_{400'})}{\sin(\alpha_{100'} + \alpha_{400'})} \tag{5.8}$$

and

$$a_2 = \frac{a_{23}\sin(\alpha_{123} + \varphi - \alpha_{300'})}{\sin(\alpha_{20'0} + \alpha_{30'0})} \tag{5.9}$$

where

$$\alpha_{100'} = \arccos\left(\frac{\bar{B}_4{}^2 - \bar{B}_1{}^2 - x_G^2}{2\bar{B}_1 x_G}\right), \tag{5.10}$$

$$\alpha_{0'04} = \arccos\left(\frac{\bar{B}_1{}^2 - \bar{B}_4{}^2 - x_G^2}{2\bar{B}_4 x_G}\right), \tag{5.11}$$

$$\alpha_{104} = \arccos\left(\frac{x_G^2 - \bar{B}_1{}^2 - \bar{B}_4{}^2}{2\bar{B}_1\bar{B}_4}\right), \tag{5.12}$$

$$\alpha_{00'3} = \arccos\left(\frac{\bar{B}_2{}^2 - \bar{B}_3{}^2 - x_G^2}{2\bar{B}_3 x_G}\right), \tag{5.13}$$

$$\alpha_{00'2} = \arccos\left(\frac{\bar{B}_3{}^2 - x_G^2 - \bar{B}_2{}^2}{2x_G\bar{B}_2}\right), \tag{5.14}$$

$$\alpha_{20'3} = \arccos\left(\frac{x_G^2 - \bar{B}_2{}^2 - \bar{B}_3{}^2}{2\bar{B}_2\bar{B}_3}\right). \tag{5.15}$$

and the variable weights $\bar{B}_i \equiv (\bar{B}_i)_{1234}$ are taken from the system of equations (5.1), (5.2), (5.3) or (5.4), (5.5) and (5.6). Taking into account Lemma 6, we express $(\bar{B}_i)_{1234}$ as a function of $\bar{B}_4)_{1234}$

By taking into account the distance of a_{12} from the line defined by A_0 and A_0', and the distance of A_2 from the line which passes through A_1 and is parallel to the line defined by A_0 and A_0', we express l as a function w.r. to $\bar{B}_4$, x_G and the five Euclidean elements a_{12}, a_{13}, a_{23}, a_{34}, a_{24}, for $i = 1, 2, 3, 4$.

$$l = a_1 \cos(\alpha_{100'}) + a_2 \cos(\alpha_{20'0}) + a_{12} \cos(\varphi) \tag{5.16}$$

By letting $l \equiv \epsilon$ a positive real number we derive a nonlinear equation which depends only on the absorbing rate of the generalized Gauss-Menger tree of degree three, where the 3wFT point A_0 remains the same because the weights $(\bar{B}_i)_{1234}$ satisfy the dynamic plasticity equations of the fixed variable weighted Fermat-Torricelli tree of degree four concerning the same boundary quadrilateral $A_1A_2A_3A_4$ where the position of A_0 remains invariant in $\mathbb{R}^2$.

By letting $l \equiv \epsilon_i$ a decreasing sequence which converge to zero, A_0' will approach the fixed position of A_0.

A universal constant u_c may occur if $x_G(\epsilon_k) \to u_c$ for every positive real value of the variable weight $(\bar{B}_4)_{1234}$. Thus, a weighted Fermat-Torricelli degree four regarding the same boundary quadrilateral $A_1A_2A_3A_4$ would exist with a constant absorbing rate and could offer an analytical solution of the 4wFT problem in $\mathbb{R}^2$ which is contradictory with the result of Theorem 4. □

The location of the 4wFT point for a convex quadrilateral or the position of a weighted Fermat-Torricelli tree of degree four is given by the following lemma which has been derived in [13] ($(B_1)_{1234} > (B_2)_{1234} > (B_3)_{1234} > (B_4)_{1234}$):

Lemma 7. [13, Formula (5),(6),(7),(8),p. 413] *The following system of equations allow us to compute the position of the 4wFT point A_0 and provides a necessary condition to locate it at the interior of the convex quadrilateral $A_1A_2A_3A_4$:*

$$\cot(\alpha_{013}) = \frac{\sin(\alpha_{213}) - \cos(\alpha_{213})\cot(\alpha_{102}) - \frac{a_{31}}{a_{12}}\cot(\alpha_{304}+\alpha_{401})}{-\cos(\alpha_{213}) - \sin(\alpha_{213})\cot(a_{102}) + \frac{a_{31}}{a_{12}}}, \tag{5.17}$$

$$\cot(\alpha_{013}) = \frac{\sin(\alpha_{314}) - \cos(\alpha_{314})\cot(\alpha_{401}) + \frac{a_{31}}{a_{41}}\cot(\alpha_{304}+\alpha_{401})}{\cos(\alpha_{314}) + \sin(\alpha_{314})\cot(\alpha_{401}) - \frac{a_{31}}{a_{41}}}, \tag{5.18}$$

$$\begin{aligned}(B_3)^2_{1234} = (B_1)^2_{1234} + (B_2)^2_{1234} + (B_4)^2_{1234} + \\ 2(B_2)_{1234}(B_4)_{1234}\cos(\alpha_{401}+\alpha_{102}) + 2(B_1)_{1234}(B_2)_{1234}\cos(\alpha_{102}) + \\ +2(B_1)_{1234}(B_4)_{1234}\cos(\alpha_{401}),\end{aligned} \tag{5.19}$$

$$\cot(\alpha_{304}+\alpha_{401}) = \frac{(B_1)_{1234} + (B_2)_{1234}\cos(\alpha_{102}) + (B_4)_{1234}\cos(\alpha_{401}))}{(B_4)_{1234}\sin(\alpha_{401}) - (B_2)_{1234}\sin(\alpha_{102})}, \tag{5.20}$$

$$\alpha_{203} = 2\pi - \alpha_{102} - \alpha_{304} - \alpha_{401}, \tag{5.21}$$

$$a_{01} = a_{14}\frac{\sin(\alpha_{013}+\alpha_{314}+\alpha_{401})}{\sin\alpha_{401}} \tag{5.22}$$

and

$$a_{04} = a_{14}\frac{\sin(\alpha_{013}+\alpha_{314})}{\sin\alpha_{401}}. \tag{5.23}$$



FIGURE 2. A fixed weighted Fermat-Torricelli tree of degree four having a universal Fermat-Torricelli minimum value $u_{FT} = \min x_G = 3.8088826$ with respect to a boundary weighted rectangle taken from Example 2.

Example 2. *Given a rectangle $A_1A_2A_3A_4$ in $\mathbb{R}^2$ such that: $A_1 = (0,0)$, $A_2 = (7,0)$, $A_3 = (7,4)$, $A_4 = (0,4)$, with side lengths $a_{12} = a_{34} = 7$, $a_{23} = a_{41} = 4$ and initial positive weights $(B_1)_{1234} = 3$, $(B_2)_{1234} = 2.5$ $(B_3)_{1234} = 1.7$ $(B_4) = 1.5$ which correspond to the vertices A_1, A_2, A_3 and A_4, respectively. By substituting these data in (5.17), (5.18) (5.19), (5.20), (5.21) of Lemma 7 we obtain the location of the 4wFT point $A_0 = (2.8274502, 1.2787811)$ which coincides with result obtained by the Weiszfeld algorithm (*[11],[12]*) with 7 digit precision and we derive that:*

$$\begin{aligned} &\alpha_{102} = 138.625^\circ, \alpha_{203} = 50.1502^\circ, \\ &\alpha_{304} = 102.986^\circ, \alpha_{401} = 68.2392^\circ. \end{aligned} \tag{5.24}$$

By substituting (5.24) in (5.1), (5.2) and (5.3) or in (5.4), (5.5) and (5.6) we obtain the dynamic plasticity equations of $A_1A_2A_3A_4$:

$$\begin{aligned} (B_1)_{1234} &= 4.2239621 - 0.8159745(B_4)_{1234}, \\ (B_2)_{1234} &= 0.8393665 + 1.1070888(B_4)_{1234}, \\ (B_3)_{1234} &= 3.6366712 - 1.2911143(B_4)_{1234}. \end{aligned} \tag{5.25}$$

Then, we replace (5.25) for $(B_4)_{1234} = 1.5$, $(B_4)_{1234} = 1.2$, $(B_4)_{1234} = 1.7$, $(B_4)_{1234} = 1.7728955$, in the objective function (3.1) of a generalized Gauss tree of degree three, taking into account (3.4), (3.6) and (3.5) from Lemma 4 and by maximizing (3.1) w.r. to x_G, we derive the following table (Fig. 2).

$(B_1)_{1234}$	$(B_2)_{1234}$	$(B_3)_{1234}$	$(B_4)_{1234}$	x_G	f
3	*2.5*	*1.7*	*1.5*	*3.8192408*	*34.5746856*
3.2447927	*2.1678731*	*2.0873328*	*1.2*	*3.8543169*	*34.6371118*
2.8368055	*2.7214176*	*1.4417756*	*1.7*	*3.8096235*	*34.5330567*
2.7773246	*2.8021194*	*1.3476592*	*1.7728955*	*3.8088826*	*34.5178864*



FIGURE 3. A fixed weighted Fermat-Torricelli tree of degree four having a universal Fermat-Torricelli minimum value $u_{FT} = \min x_G = 3.66326$ with respect to a boundary weighted rectangle taken from Example 3.

By substituting $l = 0.0000001$ *and (5.25) in (5.16) and by minimizing (5.16) w.r. to the variables* x_G *and* $(B_4)_{1234}$, *we obtain the universal minimum Fermat-Torricelli value* $u_{FT} = 3.8088826$ *for* $(B_4)_{1234} = 1.7728955$ *(see also the above Table, Fig. 2).*

The universal absorbing rate is $\frac{u_{FT}}{\sum_{i=1}^{4}(B_i)_{1234}} = \frac{3.8088826}{8.7} = 0.4378025$.

Example 3. *Given the same rectangle* $A_1A_2A_3A_4$ *in* $\mathbb{R}^2$ *with the one considered in Example 6 and initial positive weights* $(B_1)_{1234} = 3.1$, $(B_2)_{1234} = 2.3$ $(B_3)_{1234} = 1.7$ $(B_4) = 1.4$ *which correspond to the vertices* A_1, A_2, A_3 *and* A_4, *respectively. By substituting these data in (5.17), (5.18) (5.19), (5.20), (5.21) of Lemma 7 we obtain the location of the 4wFT point* $A_0 = (2.381487, 1.1855484)$ *which coincides with result obtained by the Weiszfeld algorithm (*[11],[12]*) with 7 digit precision and we derive that:*

$$\begin{aligned} &\alpha_{102} = 139.138^\circ, \alpha_{203} = 45.7542^\circ, \\ &\alpha_{304} = 98.8792^\circ, \alpha_{401} = 76.2283^\circ. \end{aligned} \tag{5.26}$$

By substituting (5.26) in (5.1), (5.2) and (5.3) or in (5.4), (5.5) and (5.6) we obtain the dynamic plasticity equations of $A_1A_2A_3A_4$:

$$\begin{aligned} (B_1)_{1234} &= 4.1823652 - 0.7731178(B_4)_{1234}, \\ (B_2)_{1234} &= 0.49794 + 1.2871855(B_4)_{1234}, \\ (B_3)_{1234} &= 3.8196947 - 1.5140677(B_4)_{1234}. \end{aligned} \tag{5.27}$$

By substituting $l = 0.0000001$ *and (5.25) in (5.16) and by minimizing (5.16) w.r. to the variables* x_G *and* $(B_4)_{1234}$, *we obtain the universal minimum Fermat-Torricelli value* $u_{FT} = 3.66326$ *for* $(B_4)_{1234} = 1.8199325$.

The universal absorbing rate is $\frac{u_{FT}}{\sum_{i=1}^{4}(B_i)_{1234}} = \frac{3.66326}{8.5} = 0.4309717$ *(Fig. 3).*

Definition 13. *A universal absorbing Fermat-Torricelli set with respect to a fixed variable weighted Fermat-Torricelli tree of degree four is the set of values of the*

Gauss variable x_G which maximizes the objective function of a generalized Gauss Menger tree of degree three which is concentrated on the same weighted Fermat-Torricelli tree of degree four, such that the variable weights satisfy the dynamic plasticity equations of the weighted boundary quadrilateral.

Definition 14. *We call a universal Fermat-Torricelli minimum value u_{FT} the minimum of the universal Fermat-Torricelli set regarding a fixed variable weighted Fermat-Torricelli tree of degree four.*

Remark 3. *The weighted Fermat-Torricelli tree of degree four taken from example 2 has greater universal Fermat-Torricelli minimum value u_{FT} and universal absorbing rate than the weighted Fermat-Torricelli tree of degree four w.r. to the same boundary quadrilateral $A_1A_2A_3A_4$ with different weights*

6. Steady trees and evolutionary trees for a boundary convex quadrilateral

Consider a universal Fermat-Torricelli set U and minimum value u_{FT} which corresponds to a fixed variable weighted Fermat-Torricelli tree of degree four regarding a boundary convex quadrilateral, which is derived as a limiting tree structure from a generalized Gauss tree of degree.

Definition 15. *A steady tree of degree four is a weighted Fermat-Torricelli tree of degree four having as storage at the 4wFT point A_0 a positive quantity less than u_{FT}.*

Definition 16. *An evolutionary tree of degree three is a generalized Gauss Menger tree of degree three which is derived by a weighted Fermat-Torricelli tree of degree four having as storage at the 4wFT point A_0 a positive quantity equal or greater than u_{FT} and then decreases by an absorbing rate $0 < a_G < u_{FT}$.*

Example 4. *Consider the same weighted Fermar-Torricelli tree four that we have obtained in Example 2 for the boundary rectangle $A_1A_2A_3A_4$. A steady tree is a weighted Fermat-Torricelli tree of degree four with storage level at A_0 less than $u_{FT} = 3.8088826$.*

When the storage quantity at A_0 reaches $u_{FT} = 3.8088826$, it stimulates the steady tree which starts to evolute (Fig. 4). Thus, the universal minimum Fermat-Torricelli value unlocks the evolution of a generalized Gauss-Menger tree which could be derived by a weighted Fermat-Torricelli tree of degree four for the same boundary rectangle (or convex quadrilateral).

For instance, if the storage level reaches $u = 3.8543169 > u_{FT} = 3.8088826$ and spends $a_G = 0.5$, then we derive an evolutionary generalized Gauss tree having $x_G = 3.3543169$ (Fig. 5).

The evolutionary Gauss-Menger tree is obtained by substituting $x_G = 3.3543169$, $(B_1)_{1234} = 3.2447927$, $(B_2)_{1234} = 2.1678731$, $(B_3)_{1234} = 2.0873328$, $(B_4)_{1234} = 1.2$ in (3.4), (3.6) and (3.5) from Lemma 4. Thus, we get:

$$a_1 = 1.6642065, a_2 = 2.7738702, a_3 = 3.6321319,$$
$$a_4 = 3.4873166, l = 3.3543169. \tag{6.1}$$

Suppose that the storage level reaches $u = 3.82 > u_{FT} = 3.8088826$ and spends $a_G = 0.2$, then we derive another evolutionary weighted Fermat-Torricelli tree



FIGURE 4. An evolutionary tree of degree four having an initial storage level 3.8543169 and absorbing rate $a_G = 0$ with respect to a boundary weighted rectangle taken from Example 2 for $(B_1)_{1234} = 3.2447927$, $(B_2)_{1234} = 2.1678731$, $(B_3)_{1234} = 2.0873328$, $(B_4)_{1234} = 1.2$.



FIGURE 5. An evolutionary tree of degree three having an initial storage level 3.8543169 and absorbing rate $a_G = 0.5$ with respect to a boundary weighted rectangle taken from Example 2 for $(B_1)_{1234} = 3.2447927$, $(B_2)_{1234} = 2.1678731$, $(B_3)_{1234} = 2.0873328$, $(B_4)_{1234} = 1.2$.

of degree four and a generalized Gauss tree of degree three having $x_G = 3.62$ *(Figs. 6, 7,8).*

By substituting $u = x_G = 3.82$ *and the dynamic plasticity equations (5.25) taken from Example 2 in (3.1), the maximum of (3.1) w.r. to* $(B_4)_{1234}$ *yields* $(B_4)_{1234} = 1.4901507$ *or* $(B_4)_{1234} = 2.0556426$. *By letting* $(B_4)_{1234} = 1.4901507$ *in (5.25), we get:* $(B_1)_{1234} = 3.0080371$, $(B_2)_{1234} = 2.4890958$, $(B_3)_{1234} = 1.7127149$,

and by letting $(B_4)_{1234} = 2.0556426$ *in (5.25), we get:* $(B_1)_{1234} = 2.5466101$, $(B_2)_{1234} = 3.1151456$, $(B_3)_{1234} = 0.9826002$.

The first evolutionary Gauss-Menger tree is obtained by substituting $x_G = 3.62$, $(B_1)_{1234} = 3.0080371$, $(B_2)_{1234} = 2.4890958$, $(B_3)_{1234} = 1.7127149$, $(B_4)_{1234} =$



FIGURE 6. An evolutionary tree of degree four having an initial storage level 3.82 and absorbing rate $a_G = 0$ with respect to a boundary weighted rectangle taken from Example 2 for $(B_1)_{1234} = 3.0080371$, $(B_2)_{1234} = 2.4890958$, $(B_3)_{1234} = 1.7127149$, $(B_4)_{1234} = 1.4901507$.



FIGURE 7. An evolutionary tree of degree three having an initial storage level 3.82 and absorbing rate $a_G = 0.2$ with respect to a boundary weighted rectangle taken from Example 2 $(B_1)_{1234} = 3.0080371$, $(B_2)_{1234} = 2.4890958$, $(B_3)_{1234} = 1.7127149$, $(B_4)_{1234} = 1.4901507$.

1.4901507 *in (3.4), (3.6) and (3.5) from Lemma 4 (Fig. 7). Thus, we get:*

$$a_1 = 2.5638686, a_2 = 3.4255328, a_3 = 4.2080591,$$
$$a_4 = 3.6397828, l = 1.5309344. \tag{6.2}$$

The second evolutionary Gauss-Menger tree is obtained by substituting $x_G = 3.62, (B_1)_{1234} = 2.5466101, (B_2)_{1234} = 3.1151456, (B_3)_{1234} = 0.9826002.$ $(B_4)_{1234} = 2.0556426.$ *in (3.4), (3.6) and (3.5) from Lemma 4 (Fig. 8). Thus, we get:*

$$a_1 = 2.6836315, a_2 = 3.0204233, a_3 = 4.0857226,$$
$$a_4 = 3.6424502, l = 1.8001622. \tag{6.3}$$



FIGURE 8. An evolutionary tree of degree three having an initial storage level 3.82 and absorbing rate $a_G = 0.2$ with respect to a boundary weighted rectangle taken from Example 2 $(B_1)_{1234} = 2.5466101$, $(B_2)_{1234} = 3.1151456$, $(B_3)_{1234} = 0.9826002$. $(B_4)_{1234} = 2.0556426$.

REFERENCES

[1] V. Boltyanski, H. Martini, V. Soltan, *Geometric Methods and Optimization Problems*, Kluwer, Dordrecht-Boston-London, 1999.

[2] D. Cieslik, Shortest connectivity. An introduction with applications in phylogeny, Springer-Verlag, New York (2005).

[3] R. Courant and H. Robbins, *What is Mathematics?* Oxford University Press., New York, 1951.

[4] E.Engelbrecht, *Planimetrischer Lehrsatz*. Arch. Math. Phys. **60**(1877), 447–448.

[5] E.N. Gilbert and H.O. Pollak, *Steiner Minimal trees*, SIAM Journal on Applied Mathematics.**16** (1968), 1-29.

[6] S. Gueron and R. Tessler, *The Fermat-Steiner problem*, Amer. Math. Monthly **109** (2002), 443-451.

[7] V. Jarnik and R. Kossler, *O minimalnich grafeth obeahujicich n danijch bodu*, Cas. Pest. Mat. a Fys., 63 (1934), 223–235.

[8] A. O. Ivanov and A. A. Tuzhilin, *Weighted minimal binary trees*. (Russian) Uspekhi Mat. Nauk 50 (1995), no. 3(303), 155–156; translation in Russian Math. Surveys 50 (1995), no. 3, 623–624.

[9] A. O. Ivanov and A. A. Tuzhilin, *Branching solutions to one-dimensional variational problems*. World Scientific Publishing Co., Inc., River Edge, NJ, 2001.

[10] Y.S. Kupitz and H. Martini, *Geometric aspects of the generalized Fermat-Torricelli problem*, Bolyai Society Mathematical Studies, **6**, (1997) , 55-127.

[11] E. Weiszfeld, *Sur le point lequel la somme des distances de n points donnes est minimum*, Tohoku Math. J. **43**, (1937), 355–386.

[12] E. Weiszfeld, *On the point for which the sum of the distances to n given points is minimum*, Ann. Oper. Res. **167** (2009), 7-41.

[13] A.N. Zachos and G. Zouzoulas, *An evolutionary structure of convex quadrilaterals*, J. Convex Anal., **15**, no. 2 (2008), 411–426.

[14] A.N. Zachos and G. Zouzoulas, *An evolutionary structure of convex quadrilaterals. Part II.*, J. Convex Anal., **20**, no. 2 (2013), 483–493.

[15] A.N. Zachos, *A weighted Steiner minimal tree for convex quadrilaterals on the two-dimensional K-plane*, J. Convex Anal., **18**, no. 1 (2011), 139–152.

[16] A.N. Zachos, *A Plasticity Principle of Convex Quadrilaterals on a Convex Surface of Bounded Specific Curvature* ,Acta. Appl. Math. (2014),**129**, no. 1, 81–134.

[17] A.N. Zachos, *Solving a generalized Gauss problem*, Mediterr. J. Math. 12 (2015), 1069-1083.

Greece
E-mail address: azachos@gmail.com